



Journal Website:
<https://theusajournals.com/index.php/ijmscr>

Copyright: Original
content from this work
may be used under the
terms of the creative
commons attributes
4.0 licence.



MYRINGOPLASTY WITH APPLICATION XENOGRFT FROM SHEEP PERICARDIA

Submission Date: April 14, 2023, Accepted Date: April 19, 2023,

Published Date: April 24, 2023

Crossref doi: <https://doi.org/10.37547/ijmscr/Volume03Issue04-13>

Matkuliev K.M.

Researcher Tashkent Medical Academy, Uzbekistan

Khasanov U.S.

Researcher Tashkent Medical Academy, Uzbekistan

Rakhimjonova G.A.

Researcher Tashkent Medical Academy, Uzbekistan

Bakieva Sh.K.

Researcher Tashkent Medical Academy, Uzbekistan

Abdullaeva N.N.

Researcher Tashkent Medical Academy, Uzbekistan

ABSTRACT

To date, there is no consensus on the question of which tissues are more appropriate to use for tympanic membrane plastic surgery. We decided to study the use of xenograft from sheep pericardium in myringoplasty in rabbits in the experiment, and in patients with chronic dry mesotympanitis. Experimental morphological studies using a xenograft were carried out in 28 rabbits. In the experiment, a positive result was obtained in 25 (89.3%) rabbits. In addition, 77 patients with dry perforation of the tympanic membrane were examined. To compare the results of the study, these patients underwent myringoplasty: 54 patients with the use of a xenograft from the sheep pericardium, 23 patients with the use of the fascia of the temporal muscle. Reliable engraftment of a xenograft from sheep pericardium after surgery was observed in 48 patients (88.9%), and with the use of temporal muscle fascia in 20 patients (86.9%).

KEYWORDS

Xenograft, myringoplasty, temporal muscle fascia, flap.

INTRODUCTION

Despite the significant progress made in reconstructive and restorative surgery of the middle ear over the past two to three decades, the problem of surgical restoration of the integrity of the tympanic membrane is still relevant. No less important is the problem of choosing an effective plastic material for the formation of the neotympanic membrane.

Most otosurgeons prefer the autofascia of the temporal muscle, which is similar in nature to the tympanic membrane and convenient to use [5,8]. However, it, like all soft tissue grafts, often atrophies, which causes a recurrence of perforation or turns into a flabby scar, which significantly reduces the functional effect of the operation [2]. For this reason, in recent years, more rigid, usually multi-layered grafts have been used, which provide better morphological results.

However, due to the use of several heterogeneous tissues, the formed neotympanic membrane differs from the natural one in its physical and acoustic properties, which inevitably affects the functional result of myringoplasty.

Thus, the current state of the problem of reconstructive and restorative surgery of the middle ear, in particular the plastics of tympanic membrane defects, dictates the urgent need for further search for both more adequate transplants and improvement of methods of surgical interventions.

Achievements in reconstructive surgery and tissue conservation have found their expression in the use of flaps - homotransplants, mainly from connective tissue [11] (pericardium, aortic valve, dura mater, sclera, cornea, etc.). The most popular is the preserved dura

mater, which has low antigenicity, good revascularization, resistance to infections, and quickly interacts with the receptive bed [3,7].

So V.I. Rodin et al.[4] using a preserved dura mater, they achieved tympanic membrane defect closure in 93% of patients, but according to M. Tos [1], the dura mater is quite rigid, dense and thick material, and this is one of its main drawbacks. Allografts were also used for myringoplasty: the dura mater of an adult and fetus [6], the tympanic membrane of an adult and fetus. Allografts can be transmitters of such dangerous infections as AIDS, hepatitis, syphilis, as well as genetic diseases.

Many years of clinical experience with the use of these grafts in surgical interventions on the middle ear revealed a number of disadvantages: rejection of the plastic material, recurrence of perforation in the neotympanic membrane, the need for additional operations to collect the graft, which negatively affects the anatomical and functional results of surgical treatment of patients. Therefore, in modern otosurgery, the development of new highly effective grafts for tympanic membrane defect repair is an urgent and important problem.

Recently, many foreign authors have used xenografts for myringoplasty [9,10]. In our republic, R.O. Mukhamadiev created a xenograft [12] from the pericardium of sheep (author's certificate No. 002-03/145 dated March 28, 2003), which was used by the author extrasclerally for various pathologies of the organ of vision. The xenograft attracts attention due to the availability and almost unlimited supply of plastic material..

The aim of our study is a clinical and experimental substantiation of the effectiveness of the use of a xenograft from the sheep pericardium in myringoplasty.

Material and methods. Experimental and morphological studies were carried out on 28 rabbits using a xenograft from sheep pericardium with dry mesotympanitis on the right ear (main group). The control group consisted of the same rabbits (left ear), in which dry mesotympanitis was modeled. Morphological features of xenograft engraftment in experimental animals we made on the 3rd, 7th, 14th, 21st days, 1 month and 3 months after the operation. In the period from 3 days to 3 months after the operation, the animals were euthanized by air embolism and subjected to post-mortem examination. The extracted xenograft was studied macro- and microscopically. The pieces were fixed in 10% neutral formalin solution. After washing in running water, dehydration was carried out in alcohol and chloroform, and then they were filled with paraffin and wax. Histological sections were stained with hematoxylin-eosin. Collagen fibers were detected by the Vann-Gieson method.

Results of the morphological study:

On the 3rd day after myringoplasty in the area and circumference of the xenograft, dyscirculatory changes were noted in the form of plethora of vessels of the microcirculatory bed, diapedetic hemorrhage in the perivascular zone, expansion of postcapillary venules, and marginal location of white blood cells. Directly in the circumference of the transplanted pericardium, these changes were accompanied by hemorrhages, loosening and fibrinoid necrosis of the outer membrane of the pericardium. The fibrous structures of the middle membranes of the pericardium were subjected to edema, loosening and moderate swelling. The above morphological and

functional changes in the soft tissues of the middle ear indicate the development of acute discirculatory, edematous-destructive changes in the surgical injury and xenograft.

On the 7th day after myringoplasty in the zone and circumference of the xenograft, dyscirculatory changes turn into response inflammatory processes in the form of hyperemia of the vessels of the microcirculatory bed, thinning of their walls and migration of both polynuclear and mononuclear leukocytes into the perivascular zones. An inflammatory infiltrate is formed on the border of the pericardium and the surrounding tissue. On the part of the xenograft, loosening and destruction of its outer shell in the form of fibrinoid necrosis, in which activated lymphoid and macrophage cells appear, was noted. In the middle membrane of the pericardium, against the background of loosening and homogenization of fibrous structures, the appearance of active cells, both of the proper pericardial and reactive origin, was noted.

On the 14th day after myringoplasty, fusion of the tissue structures of the pericardium with the surrounding soft tissue and the bone base is noted along the edges of the xenograft. At the same time, the tissue structures of the pericardium are completely destroyed and mixed with the surrounding structural elements of both soft and bone tissue. From the side of the skin of the external auditory canal, there was also a complete fusion of the pericardium with the structural elements of the skin. Only in the areas of the surgical incision, the appearance of a small proliferative infiltration, consisting of granulation tissue and inflammatory cells, is observed.

21 days after myringoplasty. There was a decrease in the volume of destructive changes in the composition of the xenograft. Maturation and differentiation of

young histiocytic cells of inflamed pericardial foci into mature histiocytes and fibrocytes were observed with the development of fibrous structures, which, merging with the fibrous elements of the pericardium, form dense and thick fibrillar structures. On the part of the bone tissue, the appearance of a thin layer similar to the periosteum, consisting of cellular-fibrous connective tissue, was determined. On the part of the skin of the external auditory canal, the formation of a mature connective tissue from the inflammatory-granulation tissue and its fusion with the fibrous tissue of the hypodermis and dermis was also noted.

One month after the experiment, the development of regressive phenomena in the composition of myringoplasty and surrounding tissues was noted in the form of the disappearance of inflammatory cells, maturation and differentiation of cambial histiocytic cells into mature histiocytes and fibrocytes. Moreover, such a regressive differentiation of connective tissue cells was accompanied by a decrease in their number. The number of thin-walled vessels also decreased with stabilization of hemodynamic and edematous phenomena. Fibrous structures in the composition of the newly formed connective tissue and pericardium also decreased in volume, turned into separate bundles, which were tightly and closely connected on one side with bone tissue, and on the other with the dermis of the skin.

Three months after pericardial myringoplasty in the surgical field, we saw the complete disappearance of all types of general pathological, inflammatory and regenerative changes. There was a complete fusion of the pericardial tissue with the surrounding tissues and microscopically, both in composition and in maturity, did not differ from the local connective tissue. Only the preservation of small areas with moderately differentiated fibrocytic cells and signs of angiomatosis was observed.

Macroscopic changes in the xenograft were assessed using an otoscope and a microscope. According to the results of experimental studies, it was noted: complete rejection of the graft 1 (3.57%) 3-day, prolapse of the graft in the tympanic cavity 1 (3.57%) 14-day, incomplete healing of the graft 1 (3.57%) 21-day.

The clinical study was carried out on the basis of the TMA clinic. The main group consisted of 54 patients who underwent myringoplasty with a xenograft from sheep pericardium. The control group consisted of 23 patients who underwent temporal muscle fascia during myringoplasty. The distribution of patients by gender was as follows: in the main group of 54 patients, there were 33 women and 21 men. In the control group, out of 23 patients, 11 were women and 12 were men.

Distribution of patients of the main and control age groups

Age	Main group (n=54)		Control group (n=23)	
	abs.	%	abs.	%
up to 19 years old	16	29,6	9	39,1
20-30 years old	25	46,3	6	26,1
31-40 years old	6	11,1	6	26,1
41-50 years old	5	9,3	-	-
51-60 years old	1	1,85	2	8,7
Over 60 years old	1	1,85	-	-

To determine the degree of hearing loss, patients with dry mesotympanitis underwent the following clinical and audiological studies: a study of hearing in whispered and colloquial speech, tuning fork studies using C128 and C2048 tuning forks. Tests were carried out - Rinne, Weber, Schwabach, Jelly, Bing and the Federici occlusion test. Audiometric studies were performed on an MA-31 audiometer manufactured by Klamann and Grahnert (Germany). The differential threshold for the perception of sound intensity (TS) was determined by the Luscher method at a frequency of 2000 Hz and at a sound intensity of 40 dB above the threshold, according to the generally accepted method. The state of patency of the Eustachian tube was also determined, which was crucial for achieving a good functional result after the operation. When performing myringoplasty, we were guided by the following indicators: all patients had central perforations, patency of the auditory tube of grade 0-I, no morphological changes in the tympanic membrane mucosa during otomicroscopy, and a positive cotton test.

In patients, hearing loss I-II degree, assessed according to the international classification, prevailed.

Audiometric indicators of auditory function were performed at different times: before surgery, the nearest period, after 6 months and after 1 year.

Before surgery, air conduction was 42.2 ± 1.6 dB, bone-air interval 29.1 ± 1.1 dB. After the operation, respectively, the air conduction was 25.1 ± 0.90 dB and the bone-air interval was 13.0 ± 0.47 dB. After the operation, the air conduction decreased by 17.1 dB and the bone-air interval by 16.1 dB.

This table shows that before the operation, air conduction was 41.2 ± 1.7 dB, bone-air interval 28.0 ± 1.1 dB. After the operation, respectively, the air conduction was 27.3 ± 1.1 dB and the bone-air interval was 14.0 ± 0.56 dB. After the operation, the air conduction decreased by 13.9 dB and the air-bone interval by 14.0 dB.

In the late period after myringoplasty, a positive result was noted - in the main group 88.9% and in the control group 86.95% of patients.

Conclusion. A positive macroscopic result of the healing of a xenograft from the pericardium of a sheep during myringoplasty was obtained in 89.3% of rabbits. When analyzing the above microscopic data, it can be noted that in the early stages in the zone and circumference of the xenograft, reactive pathomorphological changes of a protective and restorative nature develop in the form of discirculation, edema, and loosening of tissue structures. These changes in the nearest terms of the experiment turn into an inflammatory-recovery process in the form of the formation of inflammatory granulation tissue at the border of the xenograft with subsequent fusion with the tissue elements of the pericardium, and from the side of the external auditory canal it is covered with skin.

The final clinical and morphological outcomes of myringoplasty in the main group of operated patients show complete tympanic membrane closure in 88.9% of patients, and in the control group of patients in 86.95% of cases. The audiometric parameters of patients with the use of a xenograft during myringoplasty are not inferior to the fascia of the temporal muscle. Thus, the use of a xenograft from sheep pericardium in myringoplasty is an alternative, more effective method, which, at the same time, being less traumatic, reduces the length of stay of patients in the hospital.

REFERENCES

1. Filippo N. et al. Biocompatibility evaluation criteria for novel xenograft materials: distribution and quantification of remnant nucleic acid and alpha-gal epitope. *J Stem Cell Res Ther* // *J Stem Cell Res Ther S.* – 2013. – T. 6. – C. 2.
2. Ensari S. et al. The use of molded tympanic heterograft (lamb peritoneum) // *Ear, nose & throat journal.* – 1995. – T. 74. – №. 7. – C. 487-489.
3. Daly K. A. et al. Effect of the α Gal epitope on the response to small intestinal submucosa extracellular matrix in a nonhuman primate model // *Tissue Engineering Part A.* – 2009. – T. 15. – №. 12. – C. 3877-3888.
4. Meddings R. N. Evaluation of collagen vicryl mesh as a dural graft substance : дис. – The University of Manchester (United Kingdom), 1992.
5. Aggarwal R., Saeed S. R., Green K. J. M. Myringoplasty // *The Journal of Laryngology & Otology.* – 2006. – T. 120. – №. 6. – C. 429-432.
6. Darouassi Y. et al. Prognostic factors of myringoplasty: study of a 140 cases series and review of the literature // *The Pan African Medical Journal.* – 2019. – T. 33.
7. Huang J. et al. Comparative efficacy of platelet-rich plasma applied in myringoplasty: A systematic review and meta-analysis // *Plos one.* – 2021. – T. 16. – №. 1. – C. e0245968.
8. Schwam Z. G., Cosetti M. K. Endoscopic myringoplasty and type I tympanoplasty // *Otolaryngologic Clinics of North America.* – 2021. – T. 54. – №. 1. – C. 75-88.
9. Schwam Z. G., Cosetti M. K. Endoscopic myringoplasty and type I tympanoplasty // *Otolaryngologic Clinics of North America.* – 2021. – T. 54. – №. 1. – C. 75-88.
10. Lou Z. Endoscopic myringoplasty: comparison of double layer cartilage-perichondrium graft and single fascia grafting // *Journal of Otolaryngology-Head & Neck Surgery.* – 2020. – T. 49. – C. 1-9.
11. Demir E. et al. Endoscopic butterfly inlay myringoplasty for large perforations // *European Archives of Oto-Rhino-Laryngology.* – 2019. – T. 276. – C. 2791-2795.

12. Ersözlü T., Gultekin E. A comparison of the autologous platelet-rich plasma gel fat graft myringoplasty and the fat graft myringoplasty

for the closure of different sizes of tympanic membrane perforations //Ear, Nose & Throat Journal. – 2020. – T. 99. – №. 5. – C. 331-336.



OSCAR
PUBLISHING SERVICES