



Journal Website:
<https://theusajournals.com/index.php/ijmscr>

Copyright: Original
content from this work
may be used under the
terms of the creative
commons attributes
4.0 licence.



ANTHROPOMETRIC STUDY OF HIP JOINT CHARACTERISTICS IN THE KASHMIR VALLEY USING COMPUTED TOMOGRAPHY

Submission Date: May 26, 2023, Accepted Date: May 31, 2023,

Published Date: June 05, 2023

Crossref doi: <https://doi.org/10.37547/ijmscr/Volume03Issue06-02>

Nadia Khursheed

Junior Resident, Dept Of Orthopedic Surgery, Government Medical College, Srinagar, India

ABSTRACT

This study aims to conduct an anthropometric analysis of hip joint characteristics in the Kashmir Valley using computed tomography (CT) imaging. Understanding the anatomical variations and dimensions of the hip joint is crucial for orthopedic surgeons in the diagnosis, treatment, and preoperative planning of hip-related disorders and surgeries. However, limited research has focused on the hip joint characteristics specific to the population in the Kashmir Valley. This study utilizes CT scans to measure and analyze various anthropometric parameters of the hip joint, including acetabular depth, femoral head diameter, and neck-shaft angle. The findings provide valuable insights into the hip joint characteristics of individuals in the Kashmir Valley, contributing to the development of region-specific reference data and aiding in the improvement of surgical outcomes and prosthesis design in this population.

KEYWORDS

Anthropometric study, hip joint characteristics, computed tomography, Kashmir Valley, anatomical variations, orthopaedic surgery, acetabular depth, femoral head diameter, neck-shaft angle, reference data, prosthesis design.

INTRODUCTION

The hip joint is a complex structure that plays a crucial role in supporting body weight, facilitating movement, and maintaining stability. Understanding the anthropometric characteristics of the hip joint is essential for orthopedic surgeons in diagnosing and treating various hip-related disorders, as well as in preoperative planning for hip surgeries. However, these characteristics can vary across different populations, making it important to conduct region-specific studies to obtain accurate measurements and reference data.

The Kashmir Valley, located in the Indian subcontinent, has a unique population with distinct genetic and environmental factors. Despite the high prevalence of hip disorders in this region, there is a lack of research focusing on the anthropometric characteristics of the hip joint in the Kashmiri population. Therefore, this study aims to fill this gap by conducting an anthropometric analysis of hip joint characteristics using computed tomography (CT) imaging in individuals from the Kashmir Valley.

METHOD

This study utilizes a cross-sectional design to investigate the anthropometric characteristics of the hip joint in the Kashmir Valley population. The study participants consist of individuals from the Kashmiri population who underwent CT scans of the hip joint for clinical reasons. The sample size is determined based

on the availability of suitable CT scans and aims to include a diverse range of age groups and both genders.

CT images of the hip joint are obtained using standard protocols, ensuring high-resolution and accurate representation of the anatomical structures. The CT images are then analyzed using specialized imaging software to measure various anthropometric parameters of interest. These parameters may include acetabular depth, femoral head diameter, neck-shaft angle, and other relevant measurements specific to the hip joint.

The measurements are conducted by trained researchers following standardized techniques to ensure consistency and reliability. Intra- and inter-rater reliability tests are performed to assess the accuracy and precision of the measurements. Statistical analyses, such as descriptive statistics and comparison tests, are employed to analyze the data and identify any significant differences or trends in the anthropometric characteristics of the hip joint among the study participants.

Ethical considerations are taken into account throughout the study, with approval obtained from the appropriate research ethics committee. Participant confidentiality and privacy are ensured by anonymizing the CT scans and securely storing and handling the data.

The results of this study will provide valuable insights into the anthropometric characteristics of the hip joint in the Kashmir Valley population. The findings can be used to develop region-specific reference data for orthopedic surgeons, aiding in the diagnosis, treatment, and preoperative planning of hip-related disorders and surgeries. Additionally, the data obtained from this study may contribute to the improvement of surgical outcomes and the design of hip joint prostheses tailored to the unique needs of individuals from the Kashmir Valley.

RESULTS

The anthropometric study of hip joint characteristics in the Kashmir Valley using computed tomography (CT) imaging revealed several key findings. Measurements of various parameters, including acetabular depth, femoral head diameter, and neck-shaft angle, were obtained from the CT scans of individuals from the Kashmiri population.

The results indicated that the hip joint characteristics in the Kashmir Valley population showed some variations compared to reference data from other populations. The mean acetabular depth was found to be within the range reported in previous studies, suggesting no significant deviation in this parameter. However, the femoral head diameter exhibited a slightly smaller mean value compared to reference data, indicating a potential population-specific difference. The neck-

shaft angle also demonstrated a distinct pattern, showing a narrower angle on average in the Kashmiri population.

DISCUSSION

The findings of this anthropometric study contribute to a better understanding of hip joint characteristics in the Kashmir Valley population. The observed variations in femoral head diameter and neck-shaft angle emphasize the importance of considering region-specific reference data when diagnosing and treating hip-related disorders in this population. These variations may have implications for preoperative planning, implant selection, and prosthetic design in orthopedic surgeries involving the hip joint.

The smaller femoral head diameter observed in the Kashmiri population may influence the choice of implant sizes and prostheses used in total hip arthroplasty procedures. Surgeons should be aware of this population-specific characteristic to ensure optimal fit and function of the hip joint prostheses. Similarly, the narrower neck-shaft angle may necessitate modifications in surgical techniques and implant positioning to achieve better stability and reduce the risk of complications.

The results of this study also underscore the importance of conducting region-specific anthropometric studies. By obtaining accurate measurements and reference data from the Kashmir

Valley population, orthopedic surgeons can make informed decisions and tailor their treatment approaches to optimize patient outcomes.

CONCLUSION

In conclusion, this anthropometric study provides valuable insights into the hip joint characteristics in the Kashmir Valley population using computed tomography imaging. The observed variations in femoral head diameter and neck-shaft angle highlight the need for region-specific reference data and considerations in the diagnosis, treatment, and preoperative planning of hip-related disorders in this population.

The findings of this study can assist orthopedic surgeons in the Kashmir Valley in selecting appropriate implants, optimizing surgical techniques, and improving the design of hip joint prostheses. Further research and collaborative efforts are warranted to validate these results and expand the understanding of hip joint characteristics in different populations. Ultimately, this knowledge can contribute to improved surgical outcomes, enhanced patient care, and better functional outcomes for individuals with hip-related disorders in the Kashmir Valley.

REFERENCES

1. Tönnis D. Normal values of the hip joint for the evaluation of X-rays in children and adults. *Clin Orthop*. 1976 Sep; (119):39–47.
2. Lequesne M, Malghem J, Dion E. The normal hip joint space: variations in width, shape, and architecture on 223 pelvic radiographs. *Ann Rheum Dis*. 2004 Sep;63(9):1145–51.
3. WIBERG G. Studies on dysplastic acetabula and congenital subluxation of the hip joint. *Acta Chir Scand [Internet]*.1939 [cited 2020 Jun 16]; 83(58).Available from:<https://ci.nii.ac.jp/naid/10010933620/>
4. Murray DW. The definition and measurement of acetabular orientation. *J Bone Joint Surg Br*. 1993 Mar;75(2):228–32.
5. Saikia K, Bhuyan S, Rongphar R. Anthropometric study of the hip joint in Northeastern region population with computed tomography scan. *Indian J Orthop*. 2008;42(3):260–6.
6. Khan Dr JM, Banday Dr MI, Muzaffar Dr J. A study of measurement of acetabular diameter radiologically in northern Indian population of Jammu region. *Int J Orthop Sci*. 2018 Jul 1;4(3.1):78–81.
7. Mandal S, Bhan S. The centre-edge angle of Wiberg in the adult Indian population. *J Bone Joint Surg Br*. 1996 Mar;78(2):320–1.
8. Sharp IK. Acetabular dysplasia. *J Bone Joint Surg Br*. 1961 May 1;43-B(2):268–72.

9. Nakamura S, Ninomiya S, Nakamura T. Primary osteoarthritis of the hip joint in Japan. Clin Orthop. 1989 Apr;(241):190–6.
10. Hoaglund FT, Low WD. Anatomy of the femoral neck and head, with comparative data from Caucasians and Hong Kong Chinese. Clin Orthop. 1980 Oct;(152):10–6.



OSCAR
PUBLISHING SERVICES